

Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the <u>interactive fault map</u>.

unnamed fault zone (Class A) No. 1216

Last Review Date: 2000-10-24

citation for this record: Redsteer, M.H., compiler, 2000, Fault number 1216, unnamed fault zone, in Quaternary fault and fold database of the United States: U.S. Geological Survey website,

https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 02:17 PM.

State(s)	WHITE PINE COUNTY, NEVADA	
Physiographic province(s)	BASIN AND RANGE	
Reliability of location	Good Compiled at 1:100,000 scale.	
	Comments: Location based on 1:250,000-scale map of Dohrenwend and others (1992 #2480). Mapping based on photogeologic analysis of primarily 1:24,000-scale color aerial photography supplemented with 1:60,000-scale black-and-white aerial photography, transferred by inspection to 1:62,500-scale topographic maps and photographically reduced and directly transferred to 1:250,000-scale topographic maps. Subsequent mapping by photogeologic analysis of 1:58,000-nominal-scale color-infrared photography transferred directly to 1:100,000-scale topographic quadrangle maps enlarged to scale of the photographs.	
Geologic setting	This unnamed fault zone is located in the northernmost part of Railroad Valley between the White Pine Range to the east and the Pancake Range to the west.	
Length (km)	9 km.	
Average strike	N18°E	
Sense of movement	Normal	
Dip Direction	E	
Paleoseismology studies		
Geomorphic expression	The location of the fault is expressed by the linear morphology of the mountain and ridges adjacent to it. Dohrenwend and others (1992 #2480) show the faults as juxtaposing bedrock against Quaternary alluvium although no fault scarps in surficial materials have been noticed.	
Age of faulted surficial deposits	Late Quaternary, based on analysis of aerial photography (Dohrenwend and others, 1992 #2480).	
Historic		

earthquake		
prehistoric	late Quaternary (<130 ka) Comments: In general, these faults form bedrock escarpments, but Dohrenwend and others (1992 #2480) mapped scarps on late Quaternary deposits along the southern part of the fault zone.	
Recurrence interval		
Slip-rate	Less than 0.2 mm/yr	
category		
	Comments: Low slip-rate category is assigned on the basis of poor geomorphic preservation and relative inactivity of similar	
	distributed faults in the Basin and Range Province.	
Date and	2000	
Compiler(s)	Margaret Hisa Redsteer, U.S. Geological Survey	
References	#2480 Dohrenwend, J.C., Schell, B.A., and Moring, B.C., 1992,	
	Reconnaissance photogeologic map of young faults in the Ely 1°	
	by 2° quadrangle, Nevada and Utah: U.S. Geological Survey Miscellaneous Field Studies Map MF-2181, 1 sheet, scale	
	1:250,000.	

Questions or comments?

Facebook Twitter Google Email

Hazards

Design Ground MotionsSeismic Hazard Maps & Site-Specific DataFaultsScenarios EarthquakesHazardsDataEducationMonitoringResearch

Search	Search
--------	--------

HomeAbout UsContactsLegal