

# Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the [interactive fault map](#).

## Artillery Range fault, southern section (Class A) No. 2051b

Last Review Date: 2015-12-15

### Compiled in cooperation with the New Mexico Bureau of Geology & Mineral Resources

*citation for this record:* Machette, M.N., and Jochems, A.P., compilers, 2015, Fault number 2051b, Artillery Range fault, southern section, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, <https://earthquakes.usgs.gov/hazards/qfaults>, accessed 12/14/2020 02:22 PM.

#### Synopsis

**General:** This major basin-bounding fault crosses the eastern piedmont of the Organ Mountains and bounds bedrock hills (northern part of Franklin Mountains) that underlie the hydrologic divide between the Mesilla and Hueco basins, north of Anthony Gap. The mapped fault trace is entirely within Quaternary deposits; the northern section forms prominent scarps on deposits of middle to late Quaternary age, whereas the southern section forms large, older fault-line scarps on deposits of early to middle Quaternary age. No detailed studies have been performed on this fault owing to its location in a restricted military area.

	<p><b>Sections:</b> This fault has 2 sections. The division into two sections is based on apparent differences in recency of movement and geomorphic expression for the northern and southern parts of fault.</p>
<p><b>Name comments</b></p>	<p><b>General:</b> Named by Seager (1981 #968) for the location of the fault within Fort Bliss Anti-Aircraft (Artillery) Range. The fault extends from about 6.5 km south of White Sands, New Mexico (where it splays southwestward from the Organ Mountains fault [2052]), south and then west into a major embayment east of Anthony Gap. The name was extended by Kelley and Matheny (1983 #1005) southward to the northern end of the Franklin Mountains, where the fault joins the East Franklin Mountains fault [900] about 1 km north of the New Mexico-Texas state boundary.</p> <p><b>Section:</b> As defined here, this section extends from the westernmost trace of the fault in the gap between the Organ and Franklin Mountains to about 1 km north of the New Mexico-Texas state line where it joins the Franklin Mountains fault [900]. Along this section, the fault is primarily concealed beneath surficial deposits.</p> <p><b>Fault ID:</b> Referred to as fault 5 on figure 1 and table 2 of Machette (1987 #847).</p>
<p><b>County(s) and State(s)</b></p>	<p>DONA ANA COUNTY, NEW MEXICO</p>
<p><b>Physiographic province(s)</b></p>	<p>BASIN AND RANGE</p>
<p><b>Reliability of location</b></p>	<p>Good Compiled at 1:24,000 scale.</p> <p><i>Comments:</i> Original mapped trace from 1:125,000 scale mapping of Seager (1987 #627). Most of the fault is shown at 1:31,250 scale by Seager (1981 #968). The location of the fault was digitized at 1:24,000 scale using photogrammetry to accurately map its trace from these maps.</p>
<p><b>Geologic setting</b></p>	<p>The fault is part of a longer fault system that extends from the latitude of Capital Peak in northern White Sands Proving Grounds south to Juarez, Mexico. It joins the Organ Mountains fault [2052] on the north and the East Franklin Mountains fault [900]</p>

	on the south. The Artillery Range fault is different in that its mapped trace is entirely within Quaternary deposits, whereas other parts of the long system are at the eastern margin of strongly uplifted bedrock-cored ranges. The northern section of the fault forms the eastern margin of an intermediate-level structural block that is bounded on the west by the Organ Mountains fault [2052].
<b>Length (km)</b>	This section is 16 km of a total fault length of 34 km.
<b>Average strike</b>	N36°W (for section) versus N7°W (for whole fault)
<b>Sense of movement</b>	Normal
<b>Dip Direction</b>	E  <i>Comments:</i> High-angle normal fault based on drill holes in basin-fill sediment and gravity data (Seager, 1981 #968). No specific measurements have been made, but dips of 60–75° E. have been reported for the Organ Mountains fault [2052] to the north. Likewise, steep easterly dips were measured along the East Franklin Mountains fault [900] to the south.
<b>Paleoseismology studies</b>	
<b>Geomorphic expression</b>	The fault is mapped as having a nearly continuous but concealed trace beneath piedmont-slope and alluvial-fan deposits between the Organ and Franklin Mountains. No scarp heights are mentioned by Seager (1981 #968), but topographic maps of the fault show differential relief of less than 10–15 m between sediment of the early to middle Pleistocene Camp Rice Formation (on the upthrown block) and younger Quaternary deposits (on the downthrown block). The topographic expression of the fault is probably erosional, resulting in a fault-line escarpment, and may reflect considerable retreat from its actual position in the shallow subsurface.
<b>Age of faulted surficial deposits</b>	Mapping by Seager (1981 #968) and Seager and others (1987 #627) shows the fault cutting sediment of the early to middle Pleistocene Camp Rice Formation, but not offsetting younger deposits.
<b>Historic earthquake</b>	
<b>Most recent</b>	middle and late Quaternary (<750 ka)

<b>prehistoric deformation</b>	<i>Comments:</i> Timing based on mapping by Seager (1981 #968) and Seager and others (1987 #627). No detailed studies have been made owing to restricted access on the Fort Bliss Anti-Aircraft (Artillery) Range.
<b>Recurrence interval</b>	
<b>Slip-rate category</b>	Less than 0.2 mm/yr  <i>Comments:</i> A low slip rate is based on 10–15 m of differential relief (estimated from topographic maps) between the piedmont slope facies of the Camp Rice Formation (early to middle Pleistocene) and unfaulted deposits of early (?) late Pleistocene age (100–130 ka).
<b>Date and Compiler(s)</b>	2015 Michael N. Machette, U.S. Geological Survey, Retired Andrew P. Jochems, New Mexico Bureau of Geology & Mineral Resources
<b>References</b>	#1005 Kelley, S., and Matheny, J.P., 1983, Geology of Anthony quadrangle, Doña Ana County, New Mexico: New Mexico Bureau of Mines and Mineral Resources Geologic Map 54, 1 sheet, scale 1:24,000.  #847 Machette, M.N., 1987, Preliminary assessment of paleoseismicity at White Sands Missile Range, southern New Mexico—Evidence for recency of faulting, fault segmentation, and repeat intervals for major earthquakes in the region: U.S. Geological Survey Open-File Report 87-444, 46 p.  #968 Seager, W.R., 1981, Geology of Organ Mountains and southern San Andres Mountains, New Mexico: New Mexico Bureau of Mines and Mineral Resources Memoir 36, 97 p., 4 pls.  #627 Seager, W.R., Hawley, J.W., Kottowski, F.E., and Kelley, S.A., 1987, Geology of east half of Las Cruces and northeast El Paso 1° x 2° sheets, New Mexico: New Mexico Bureau of Mines and Mineral Resources Geologic Map 57, 3 sheets, scale 1:125,000.

[Questions or comments?](#)

[Facebook](#) [Twitter](#) [Google](#) [Email](#)

[Hazards](#)

[Design Ground Motions](#) [Seismic Hazard Maps & Site-Specific Data](#) [Faults](#) [Scenarios](#)

[Earthquakes](#) [Hazards](#) [Data](#) [Education](#) [Monitoring](#) [Research](#)

[Home](#) [About Us](#) [Contacts](#) [Legal](#)