Jump to Navigation

Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the interactive fault map.

Boney Mountain fault (Class A) No. 273

Last Review Date: 2017-07-01

citation for this record: , compiler, 2017, Fault number 273, Boney Mountain fault, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 02:52 PM.

Synopsis	
Name comments	**Fault ID:** Refers to fault number 351 of Jennings (1994).
County(s) and State(s)	CALIFORNIA
Physiographic province(s)	
Reliability of location	Compiled at 1:100,000 scale. *Comments:*
Geologic setting	
Length (km)	km.

Average strike	
Sense of movement	
Dip	
Paleoseismology studies	
Geomorphic expression	
Age of faulted surficial deposits	
Historic earthquake	
Most recent prehistoric deformation	undifferentiated Quaternary (<1.6 Ma) *Comments:*
Recurrence interval	
Slip-rate category	Unspecified
Date and Compiler(s)	2017
References	#2878 Jennings, C.W., 1994, Fault activity map of California and adjacent areas, with locations of recent volcanic eruptions: California Division of Mines and Geology Geologic Data Map 6, 92 p., 2 pls., scale 1:750,000. #8382 Yerkes, R.F., and Campbell, R.H., 2005, Preliminary geologic map of the Los Angeles 30' x 60' quadrangle, southern California: U.S. Geological Survey Open-File Report 05-1019, sheet 1, scale 1:100,000.

Search... Search

Home About Us Contacts Legal