Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the <u>interactive fault map</u>.

Baxter Spring fault zone (Class A) No. 447

Last Review Date: 2017-07-01

citation for this record: , compiler, 2017, Fault number 447, Baxter Spring fault zone, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 03:10 PM.

Synopsis	
Name comments	
County(s) and State(s)	CALIFORNIA
Physiographic province(s)	
Reliability of location	Compiled at 1:62,500 scale. Comments:
Geologic setting	
Length (km)	km.

Average strike		
Sense of movement	Normal	
Dip		
Paleoseismology studies		
Geomorphic expression		
Age of faulted surficial deposits		
Historic earthquake		
Most recent prehistoric deformation	undifferentiated Quaternary (<1.6 Ma) Comments:	
Recurrence interval		
Slip-rate category	Unspecified	
Date and Compiler(s)	2017	
References	#5577 Bailey, R.A., 1989, Geologic map of the Long Valley caldera, Mono-Inyo craters volcanic chain and vicinity, eastern California: U.S. Geological Survey Miscellaneous Investigations Series Map I-1933, scale 1:62,500.	
	#2878 Jennings, C.W., 1994, Fault activity map of California and adjacent areas, with locations of recent volcanic eruptions: California Division of Mines and Geology Geologic Data Map 6, 92 p., 2 pls., scale 1:750,000.	

Questions or comments?

Facebook Twitter Google Email

<u>Hazards</u>

<u>Design Ground MotionsSeismic Hazard Maps & Site-Specific DataFaultsScenarios</u> <u>EarthquakesHazardsDataEducationMonitoringResearch</u>

*	
Search	Search

HomeAbout UsContactsLegal