Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the <u>interactive fault map</u>.

Bitter Springs fault (Class A) No. 477

Last Review Date: 2017-07-01

citation for this record: , compiler, 2017, Fault number 477, Bitter Springs fault, in Quaternary fault and fold database of the United States: U.S. Geological Survey website, https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 03:07 PM.

Synopsis	
Name comments	
County(s) and State(s)	CALIFORNIA
Physiographic province(s)	
Reliability of location	Compiled at 1:125,000 scale. Comments:
Geologic setting	
Length (km)	km.

Average strike	
Sense of movement	
Dip	
Paleoseismology studies	
Geomorphic expression	
Age of faulted surficial deposits	
Historic earthquake	
Most recent prehistoric deformation	undifferentiated Quaternary (<1.6 Ma) Comments:
Recurrence interval	
Slip-rate category	Unspecified
Date and Compiler(s)	2017
References	#8030 Byers, F.M., Jr., 1960, Geology of the Alvord Mountain quadrangle, San Bernardino County, California: U. S. Geological Survey Bulletin 1089-A, 71 p., plate 1, map scale 1:62,500. #2878 Jennings, C.W., 1994, Fault activity map of California and adjacent areas, with locations of recent volcanic eruptions: California Division of Mines and Geology Geologic Data Map 6,
	92 p., 2 pls., scale 1:750,000.

Questions or comments?

Facebook Twitter Google Email

Hazards

<u>Design Ground MotionsSeismic Hazard Maps & Site-Specific DataFaultsScenarios</u> <u>EarthquakesHazardsDataEducationMonitoringResearch</u>

*	
Search	Search

HomeAbout UsContactsLegal