Quaternary Fault and Fold Database of the United States

As of January 12, 2017, the USGS maintains a limited number of metadata fields that characterize the Quaternary faults and folds of the United States. For the most up-to-date information, please refer to the <u>interactive fault map</u>.

Huckleberry Meadows fault (Class A) No. 783

Last Review Date: 2017-07-01

citation for this record: Bryant, W.A., compiler, 2017, Fault number 783, Huckleberry Meadows fault, in Quaternary fault and fold database of the United States: U.S. Geological Survey website,

https://earthquakes.usgs.gov/hazards/qfaults, accessed 12/14/2020 03:17 PM.

Synopsis	
Name comments	
County(s) and State(s)	SHASTA COUNTY, CALIFORNIA
Physiographic province(s)	CASCADE-SIERRA MOUNTAINS
Reliability of location	Compiled at 1: scale.
	<i>Comments:</i> Location of fault from Qt_flt_ver_3- 0_Final_WGS84_polyline.shp (Bryant, W.A., written communication to K.Haller, August 15, 2017) attributed to Page and Sawyer (2004) mapped at unspecified scale.

Geologic setting				
Length (km)	17 km.			
Average strike				
Sense of movement	Unspecified			
Dip				
Paleoseismology studies				
Geomorphic expression				
Age of faulted surficial deposits				
Historic earthquake				
Most recent prehistoric deformation	undifferentiated Quaternary (<1.6 Ma) Comments:			
Recurrence interval				
Slip-rate category	Insufficient data			
Date and Compiler(s)	2017 William A. Bryant, California Geological Survey			
References	#8225 Page, W.D., and Sawyer, T.L., 2004, Overview of late Cenozoic faulting in the Sierra Nevada Foothills (including a reassessment of faults near New Bullards Bar Dam): Unpublished Report for Pacific Gas and Electric, 28 p.			

Questions or comments?

Facebook Twitter Google Email

Hazards

Design Ground MotionsSeismic Hazard Maps & Site-Specific DataFaultsScenarios EarthquakesHazardsDataEducationMonitoringResearch

		-		
Se	-a	rc	h	

Search

-

Search... Search... Search...