

Preliminary Campbell-Bozorgnia NGA-West2 GMPE for USGS Evaluation



Kenneth W. Campbell, Ph.D.

EQECAT Inc., Beaverton, Oregon

Oregon State University, Corvallis, Oregon

Yousef Bozorgnia, Ph.D.

PEER, University of California, Berkeley, California

USGS Workshop on GMPEs, Berkeley, California

December 12–13, 2012

Contents

- Data selection criteria
- Changes from 2008 NGA GMPE
- Distribution of residuals
- Comparison with 2008 NGA GMPE
- Work to be completed

Data Selection Criteria



Data Selection Criteria



■ Earthquakes

- **M** = 3.0–5.5 California events
- **M** = 5.5–7.9 California and global events
- Known focal mechanism or fault type
- Class 1 events using 10 km CR_{JB} criteria

■ Sites


- Free field (shelters, non-embedded bldgs.)
- V_{S30} known or estimated (via proxies)
- $R_{RUP} \leq 80$ km (geometric attenuation only)
- $N \geq 5$ (**M** < 5.5), $N \geq 3$ ($5.5 \leq \mathbf{M} < 6.5$), $N \geq 1$ (**M** ≥ 6.5)

CB08 vs. CB12 Databases



CB08 vs. CB12 Databases

- CB12 data
 - $N_{eq} = 280$
 - $N_{rec} = 6525$
- CB08 data
 - $N_{eq} = 64$
 - $N_{rec} = 1561$



Changes from 2008 NGA-West1 GMPE




Changes from 2008 GMPE

- Quadralinear magnitude scaling term
 - Additional hinge at $M=4.5$
 - No longer overpredicts small M
- Hanging-Wall term from simulations
 - Functional form from Donahue study
 - Function of R_x , rupture dip and width
 - Peaks over bottom edge of fault
 - CB08 distance filter off rupture plane

Changes from 2008 GMPE

- Hypocentral depth term
 - Ground motion increases for $H_{\text{HYP}} > 7$ km
 - Statistically preferred over Z_{TOR}
 - Can use $H_{\text{HYP}} = f(Z_{\text{TOR}})$ proxy (next slide)
- Fault mechanism term
 - No longer depth-dependent
 - Goes away at small magnitudes
- Rupture plane dip term
 - Ground motion increases with dip
 - Goes away at large magnitudes

Changes from 2008 GMPE



Courtesy: Abrahamson, Chiou, Kishida, Bozorgnia

Changes from 2008 GMPE

- Shallow site-response (V_{S30}) term
 - Retained Walling et al. nonlinear model
 - Different linear dependence in Japan
 - Japan model bilinear (hinge at 200 m/sec)
- Sediment-depth ($Z_{2.5}$) term
 - California: SCEC CVM-S4, SFBA CVM-08.3.0
 - Japan: NIED Velocity Model
 - Retained Day basin model for $Z_{2.5} > 3$ km
 - Deep basin term same for Calif. and Japan
 - Shallow ($Z_{2.5} < 1$ km) term differs in Japan


Changes from 2008 GMPE

- Standard deviations
 - Similar to CB08 for $\mathbf{M} \geq 5.5$
 - Larger for $\mathbf{M} < 5.5$


Distribution of Residuals




Between-Event vs. Magnitude




Between-Event vs. Depth




Between-Event vs. Dip




Within-Event vs. Magnitude




Within-Event vs. R_{RUP} (All **M**)




Within-Event vs. R_{RUP} ($M > 6.5$)




Within-Event vs. HW Term



Within-Event vs. V_{S30}




Within-Event vs. $Z_{2.5}$




Comparison With 2008 NGA-West1 GMPE




PGA, SS, Dip=90, $V_{S30}=760$




$T=0.2\text{s}$, SS, Dip=90, $V_{S30}=760$




$T=1\text{s}$, SS, Dip=90, $V_{S30}=760$




PGA, RV, Dip=45, $V_{S30}=760$



$T=0.2s$, RV, Dip=45, $V_{S30}=760$



$T=1s, RV, Dip=45, V_{S30}=760$



Work to be Completed by Project End



Work in Progress

- Evaluate and include effects of directivity
- Evaluate new nonlinear site term
- Add regional anelastic attenuation terms
- Magnitude-dependent standard deviation
- Develop vertical GMPE